



ORIGINAL ARTICLE

Higher frequency of circulating V δ 1 $\gamma\delta$ T cells in patients with advanced schistosomiasis

Li Zheng¹  | Lixia Wang² | Yuan Hu^{3,4} | Jia Yi² | Lun Wan² | Yujuan Shen^{3,4} | Si Liu² | Xiaorong Zhou² | Jianping Cao^{3,4} 

¹Department of Immunology, Key Laboratory of Tropical Translational Medicine of Ministry of Education, Hainan Medical University-The University of Hong Kong Joint Laboratory of Tropical Infectious Diseases, Hainan Medical University, Haikou, China

²Hubei Provincial Academy of Preventive Medicine, Hubei Provincial Key Laboratory for Applied Toxicology, Hubei Provincial Center for Disease Control and Prevention, Wuhan, China

³National Institute of Parasitic Diseases, Chinese Center for Disease Control and Prevention (Chinese Center for Tropical Diseases Research), Key Laboratory of Parasite and Vector Biology, National Health Commission of People's Republic of China, WHO Collaborating Center for Tropical Diseases, Shanghai, China

⁴The School of Global Health, Chinese Center for Tropical Diseases Research, Shanghai Jiao Tong University School of Medicine, Shanghai, China

Correspondence

Jianping Cao, National Institute of Parasitic Diseases, Chinese Center for Disease Control and Prevention (Chinese Center for Tropical Diseases Research), No. 207 Rui Jin Er Road, Shanghai 200025, China.

Xiaorong Zhou. Hubei Provincial Center for Disease Control and Prevention, Wuhan 430079, China. No.35 Zhuo Dao Quan Northern Road, Wuhan 430079, China.

Email: caojpcdc@163.com(JC); 553705790@qq.com(XZ)

Funding information

This work was supported by grants from the National Natural Science Foundation of China (Nos. 8177225 and 81971969 to JC), the Hubei Provincial Natural Science Foundation of China (No. 2018CFC897 to XZ), the Hubei Province health and family planning scientific research project (No. WJ2019X004 to LZ), and NHC Laboratory of Parasite and Vector Biology (No. WSBKFKT-201709 to LZ).

Abstract

Gamma-delta ($\gamma\delta$) T cells are the bridge between natural and adaptive immunity. In the present study, peripheral blood was collected from 13 patients with advanced schistosomiasis (schistosomiasis group) and 13 uninfected people (control group) to investigate the $\gamma\delta$ T cells and their subtypes in human schistosomiasis. Compared with the control group, the proportion of V δ 1 cells and CD27⁺V δ 1⁺ cells in the schistosomiasis group increased significantly, while CD27⁻ cells and CD27⁻V δ 1⁻ cells decreased. Only the level of IL-17A differed between the groups, being significantly decreased in the schistosomiasis group. In the schistosomiasis group, there were no correlations between the liver fibrosis and subsets of $\gamma\delta$ T cells, or the level of cytokines. Additionally, the level of IL-17A correlated positively with the proportion of CD27⁻V δ 1⁻ cells. Thus, there was a higher frequency of circulating V δ 1 $\gamma\delta$ T cells in patients with advanced schistosomiasis. The decreased IL-17A might be related to the reduction in CD27⁻V δ 1⁻ cell.

KEYWORDS

fibrosis, IL-17, *Schistosoma japonicum*, schistosomiasis, $\gamma\delta$ T cell

1 | INTRODUCTION

Schistosomiasis affects more than 200 million people worldwide.¹ Previously, *Schistosoma japonicum* was endemic in China, especially in areas with many lakes, such as the 'Four Lakes Area' in Hubei province. Many patients with chronic schistosomiasis develop advanced schistosomiasis each year, despite not coming into contact with the

cercariae for a long time. For more than 50 years, researchers worldwide have studied changes in the host's immune environment during the progress of infection. While the functions of helper T (Th) cells, cytotoxic T (Tc) cells and B cells are well known, the responses and functions of $\gamma\delta$ T cells during schistosomiasis remain unclear.

$\gamma\delta$ T cells are the first line of immune defence, playing an important role in cancer, autoimmunity and infectious diseases. They

This is an open access article under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.

© 2021 The Authors. *Parasite Immunology* published by John Wiley & Sons Ltd

possess strong antimicrobial activity against viral, bacterial and other pathogens. In addition to $\gamma\delta$ T cell receptors (TCRs), $\gamma\delta$ T cells express all T-lineage-specific genes, including those encoding cell surface receptors, signalling factors, cytokines and transcription factors. Their phenotypes and functions are similar to those of activated $\alpha\beta$ T cells.² CD27 is a marker that can distinguish between the interferon- γ (IFN- γ) and interleukin-17 (IL-17) producing $\gamma\delta$ T cell subsets in a mouse model³; however, it is unknown whether the marker is applicable in humans.

Human $\gamma\delta$ T cells are divided into three main categories depending on the δ chain: V δ 1 T cells, V δ 2 T cells and V δ 3 T cells.⁴ The V δ 1 chain-expressing cells are mainly present in the epithelium of the mucous membrane, where they are involved in maintaining the integrity of epithelial tissue when faced with damage, infection and transplantation.⁵ V δ 1 T cells also appear in the peripheral blood; however, the mucosal and peripheral blood $\gamma\delta$ T cells appear to be distinct populations.⁶ V δ 1 T cells produce abundant IFN- γ and are thought to be an important source of IL-17.⁷ Most circulating $\gamma\delta$ T cells in healthy adults are V δ 2 chain-expressing T cells, with a 50–90% ratio. V δ 2 T cells are almost entirely V γ 9-expressing cells.⁸ V δ 2 T cells produce IFN- γ , IL-17A and tumour necrosis factor-alpha (TNF- α) to promote inflammation and induce anti-infective immunity in different settings of infectious diseases.⁹ Although the third class, V δ 3 T cells, represent only about 0.2% of circulating T cells, they are abundant in the liver and are increased in patients with leukaemia and in some patients with chronic viral infections. When activated, they can kill CD1d⁺ target cells and induce Th1, Th2 and Th17 cells to release cytokines.¹⁰

Most studies have observed the characteristics and explored the functions of $\gamma\delta$ T cells using mouse models, while few studies have investigated their function in humans infected with *S japonicum*.^{11–13} In 2014, Schwartz et al¹⁴ reported the similarities and differences in the characteristics of $\gamma\delta$ T cells during *S hematobium* or *S mansoni* infections in humans. However, schistosomiasis caused by *S japonicum* is a much more serious disease than that caused by other *Schistosoma* species.¹⁵ When the eggs are deposited in the liver, intestines and spleen of a patient, they can persist for a long time, leading to chronic injury and liver fibrosis. If the patient is subjected to repeated infection, even if treated, 5–10 years later they would develop advanced schistosomiasis, including cirrhosis ascites, portal hypertension and a significantly reduced quality of life.¹⁶ Therefore, it is necessary to explore the details of the change of the immune phenotype and quantity of $\gamma\delta$ T cells in patients with *S japonicum* infection.

In previous research, our team observed the characteristics of $\gamma\delta$ T cells and related cytokines in a mouse model of *S japonicum* infection, especially the subset of $\gamma\delta$ T cells that produced cytokines (IL-17/IFN- γ) in uninfected conditions, but expressed decreased levels as the infection progressed, with almost none being produced in the late stage.¹⁷ We found that the V γ 2 subset of $\gamma\delta$ T cells might play an important role in accelerating liver fibrosis by recruiting neutrophils during *S japonicum* infection. However, Hammerich et al¹⁸ found that $\gamma\delta$ T cells could protect the liver from excessive inflammation and fibrosis by

inhibiting hepatic stellate cells (HSCs). Meanwhile, Markovits et al observed anti-fibrotic characteristics of V γ 9⁺ $\gamma\delta$ T cells in systemic sclerosis,¹⁹ which presents a paradox as to whether $\gamma\delta$ T cells played a pro-fibrosis or anti-fibrosis role in the human liver; therefore, we were interested in defining the exact relationship between $\gamma\delta$ T cells and liver fibrosis in human patients infected with *S japonicum*.

The present study aimed to investigate the immune status of a population with schistosomiasis and to verify that $\gamma\delta$ T cells produce cytokines such as IL-17 during the progress of *S japonicum* infection. In addition, we explored which subtypes of $\gamma\delta$ T cells function during infection, and the relationship between $\gamma\delta$ T cells and the degree of liver fibrosis.

2 | METHODS

2.1 | Ethics approval and consent to participate

The present study was approved by the Ethics Committee of the National Institute of Parasitic Diseases, Chinese Center for Disease Control and Prevention, China (Permit Number: IPD 2017-006). Before sample collection, the survey and the procedure for sample collection were explained to the patients and the control subjects in the hospital. Each participant provided signed informed consent. All patients were treated without charge.

2.2 | Patients and sample collection

According to the 'Schistosomiasis Diagnostic Standard' (WS261-2006) provided by the Ministry of Health of the People's Republic of China, advanced schistosomiasis was defined as a patient with schistosomiasis causing liver fibrosis, portal hypertension symptoms, severe growth disorders or colon granuloma growth. Repeated or large-scale infection by *Schistosoma japonicum*, if not treated thoroughly and timely, a patient could develop advanced schistosomiasis, usually after 2–10 years of pathological development. Clinical symptoms of advanced schistosomiasis include abdominal fluid, spleen enlargement, high blood pressure, gastroesophageal varicose bleeding, large intestine granuloma lesions and severe growth retardation. People with advanced schistosomiasis were documented in the CDC (Center for Disease Control) surveillance system from the time they were first infected.

All candidates were from the Schistosomiasis control hospital of Qian-jiang City in Hubei province (China). Thirteen patients (aged 63 ± 9.63 years) with advanced schistosomiasis were selected as the schistosomiasis group, while 13 healthy controls (aged 57 ± 8.05 years) comprised the control group. All the recruited patients had no active infection. They tested negative for active eggs via the faecal miracidia hatching method. When the patients were in a stable condition, they all received praziquantel at the appropriate time. The healthy people in the control group were recruited from the physical examination centre in the same hospital. All candidates

completed forms agreeing to participate in the study. Peripheral blood samples were obtained from the two groups.

Peripheral blood from the candidates was collected into tubes with and without sodium heparin, separately. The serum samples were isolated after 2 hours of storage at room temperature. Peripheral blood mononuclear cells (PBMCs) were isolated using Ficoll-Hypaque (TBD).

2.3 | Flow cytometry

Blood samples were collected in tubes containing sodium heparin and treated with NH_4Cl erythrocyte lysis buffer. The obtained single-cell suspensions were stained with directly conjugated antibodies (listed below) for 30 minutes at 4°C in the dark in phosphate-buffered saline/1% bovine serum albumin. Dead cells were excluded by adding Fixable viability dye eFluor 780 (1:1,000; eBioscience). All antibodies were purchased from Life Technologies (Gaithersburg), except for the antibodies against V δ 1, which were from Abcam (Cambridge), and the $\gamma\delta$ TCR and CD27 antibodies, which were from eBioscience. The following antibodies were used in the $\gamma\delta$ T cell phenotyping panel: CD45-eFluor 450 (1:100; clone HI30), V δ 1-fluorescein isothiocyanate (FITC) (1:200; clone TS8.2), $\gamma\delta$ TCR-phycoerythrin (PE) (1:200; clone B1.1), CD27-Allorhycocyanin (APC) (1:100; clone O323), CD4-APC-eFluor 780 (1:200; clone OKT4), CD19- APC-eFluor 780 (1:200; clone ICRF44), CD11b-APC-eFluor 780 (1:200; clone HIB19) and fixable viability dye eFluor 780. All experiments were performed using a CytoFLEX flow cytometer (Beckman Coulter, Indianapolis, IN, USA) using CytExpert software (Beckman Coulter). Data analyses used FlowJo Software version 9.7.1 (Becton Dickinson).

2.4 | Enzyme-linked immunosorbent assay (Elisa) and serum analysis

ELISA was performed according to the manufacturer's instructions to detect the serum levels of IL-4, IL-6, IL-10, IL-17A, IL-21, transforming growth factor-beta (TGF- β), IL-1 β and IL-22. All ELISA kits were purchased from eBioscience (Invitrogen, Waltham, MA, USA).

The levels of type III procollagen (PC-III) and hyaluronic acid (HA) in the serum were assayed via a Chemiluminescent immunoassay by the Kindstar Global Department of Pathology.

2.5 | Quantitative real-time reverse transcription PCR

RNA was extracted from peripheral blood according to the manufacturer's instructions using an X-press Blood RNA kit (Omega Bio-Tek). Reverse transcriptase (RT) reactions for cDNA synthesis were performed using PrimeScript RT Master Mix (Takara). Relative mRNA expression levels were determined using quantitative real-time

polymerase chain reaction (qPCR) with an SYBR Green I PCR Master Mix kit (Takara) on an ABI ViiATM7 machine (Applied Biosystems) according to the manufacturer's protocol. The primer sequences were IL-17A forward 5'-TCCCACGAAATCCAGGATGC-3',

reverse 5'-GGATGTTTCAGGTTGACCATCAC-3'; Glyceraldehyde-3-phosphate dehydrogenase (GAPDH; control) forward 5'-CCAAGGAGTAAGACCCCTGG-3',

reverse 5'- TGGTTGAGCACAGGGTACTT-3'.

2.6 | Statistical analysis

The results were analysed using GraphPad Prism 5.0 (GraphPad Software, Inc.). All data are presented as mean \pm standard deviation. The differences between two groups were analysed using an independent t-test or Welch's t-test. The correlation between groups was tested using Spearman's rank correlation. $P < .05$ was chosen as a sensitive level of significance.

3 | Results

3.1 | The proportion of subsets of total $\gamma\delta$ T cells detected by flow cytometry

The results showed that, compared with those in the control group, the proportion of V δ 1 cells in the patients with advanced schistosomiasis increased significantly ($P = .0007$), with a significant increase in the proportion of CD27⁺V δ 1⁺ cells ($P = .0182$). We also observed a significant decrease in the ratio of CD27⁻ cells ($P = .0055$), among which the proportion of CD27⁻V δ 1⁻ cells also decreased ($P < .0001$) (Figure 1A,B).

Among the total $\gamma\delta$ T cells, the percentage of V δ 1⁺ cells was $20.53 \pm 3.05\%$ in the control group and $52.69 \pm 7.74\%$ in the schistosomiasis group, while the percentage of CD27⁺V δ 1⁺ cells was $5.46 \pm 3.79\%$ in the control group, which increased to $20.20 \pm 4.42\%$ in the schistosomiasis group. By contrast, the percentage of CD27⁻ cells was $85.54 \pm 3.85\%$ in the control group and $61.94 \pm 6.70\%$ in the schistosomiasis group, while the percentage of CD27⁻V δ 1⁻ cells was $68.18 \pm 3.17\%$ in the control group, which decreased to $29.46 \pm 5.81\%$ in the schistosomiasis group. All the results are shown as the mean \pm SEM%.

3.2 | The level of cytokines in serum detected by ELISA

The results showed that there were no significant differences between the two groups for most of the cytokines (IL-4, IL-6, IL-10, IL-21, TGF- β , IL-1 β , and IL-22); however, there was a significant decrease in the levels of IL-17A ($P = .013$), from 32.16 ± 2.55 pg/mL in the control group to 23.42 ± 1.99 pg/mL in the serum of patients with advanced schistosomiasis (Figure 1C).




FIGURE 1 Changes to the subsets of $\gamma\delta$ T cells and the related cytokines in the serum from different groups. (A, B) Representative flow cytometry plots are shown to describe the changes in the proportions of subsets of $\gamma\delta$ T cells in subjects with or without advanced schistosomiasis. Summary graphs showing the increase and decrease in subsets of $\gamma\delta$ T cells as a percentage of total $\gamma\delta$ T cells compared with those in healthy controls. (C) Differences in the cytokine profiles between the two groups. (D) The mRNA expression of *IL17A* from white cells of peripheral blood as assessed using qRT-PCR. Fold-change is the logarithmic ratio (log₂ ratio) compared with the expression level in the control. (E) The level of type III procollagen (PC-III) and hyaluronic acid (HA) in the serum were tested using a chemiluminescent immunoassay (CLIA). Data are shown as the mean \pm SE, and the *P* values are shown between the data in each column (by Student's *t*-test or with Welch's correction). *:*P* < .05. **: *P* < .01

3.3 | The level of cytokines from PBMCs tested by QRT-PCR

To check whether the level of cytokines from the PBMCs was similar to the trends shown in ELISA results, we tested the fold-change of mRNA from the same samples using qRT-PCR. The results showed that the relative mRNA level of *IL-17A* from the peripheral blood of the patients with advanced schistosomiasis was significantly reduced compared with that in the control group (*P* = .0039) (Figure 1D). By measuring the relative change of expression of *IL-17A* by qRT-PCR, we confirmed that the source of PBMCs was reduced compared with that in the healthy controls.

3.4 | The levels of HA and PC-III IN serum directly reflected the degree of hepatic fibrosis

The results showed a significant increase in HA and PC-III in the serum of the advanced schistosomiasis group compared with that in the healthy control group (Figure 1E). The HA level was 79.22 ± 7.41 ng/mL in the healthy controls, which increased to 281.70 ± 46.04 ng/mL in the schistosomiasis group (*P* = .0005). The level of PC-III was 17.30 ± 1.04 ng/mL in the healthy controls, which increased to 42.40 ± 7.28 ng/mL in the *S japonicum* group (*P* = .0026). The levels of HA and PC-III in serum directly reflected the degree of hepatic fibrosis.

3.5 | The relations between the V δ 1⁺ cells, CD27⁺V δ 1⁺ cells and cytokines

The proportions of V δ 1⁺ cells and CD27⁺V δ 1⁺ cells increased; therefore, we attempted to determine if there was a relationship between the two subsets and cytokines. The results showed that the levels of cytokines did not correlate positively with the proportion of V δ 1⁺ cells (Figure 2A) or CD27⁺V δ 1⁺ cells (Figure 2B) in total $\gamma\delta$ T cells. It was worth noting that the *P* values of IL-6 and IL-21 were .0607 and .0640 for the correlation with CD27⁺V δ 1⁺ cells, respectively, which were close to statistical significance.

3.6 | The proportions of subsets or cytokines did not correlate with liver fibrosis

In the group of patients with advanced schistosomiasis, neither the proportion of V δ 1 cells nor CD27⁺ cells correlated significantly with

liver fibrosis (Figure 3A). In addition, we detected the level of the cytokines in the serum and assessed their relationship with liver fibrosis. The results showed no significant correlation between IL-4, IL-6, IL-10, IL-17A, IL-21, TGF- β , IL-1 β , or IL-22 and liver fibrosis (Figure 3B).

3.7 | Levels of IL-17A in serum correlated with the proportions of CD27⁺V δ 1⁺ $\gamma\delta$ T cells

The results showed that the level of IL-17A in the serum correlated positively with the proportion of CD27⁺V δ 1⁺ cells among total $\gamma\delta$ T cells (*r* = 0.6703, *P* = .0122) (Figure 4). None of the other subtypes of $\gamma\delta$ T cells correlated significantly with IL-17A levels.

4 | DISCUSSION

In mouse models, $\gamma\delta$ T cells were mainly divided into V γ 1 and V γ 2 cells according to the different γ chains. In our previous study, compared with normal mice, the ratio of V γ 1 and V γ 2 cells did not change significantly in early or late infection of *S japonicum*.¹⁷ In contrast, according to the different δ chains, in human peripheral blood, $\gamma\delta$ T cells were mainly divided into V δ 1 and V δ 2 cells.²⁰ In the present study, there was a change in the percentage of V δ 1 and V δ 2 cells in patients with advanced schistosomiasis compared with that in the control group, that is, the proportion of V δ 1 cells increased, and the proportion of V δ 2 cells decreased. There might be two reasons for the increase in V δ 1 cells. The first was antigen stimulation, and the second was the relative decline of V δ 2 cells. V δ 1 cells are capable of proliferating in response to signals through TCRs and IL-15.²¹ However, during the advanced stage of schistosomiasis, the immune environment is suppressed. In the peripheral blood of healthy adults, the proportion of the V δ 2 subset among total $\gamma\delta$ T cells was greater than 70%. This subtype plays an important anti-infection and anti-tumour role. Therefore, we believe that the second explanation is more likely.

In our previous work, we demonstrated that the $\gamma\delta$ T cells lost their ability to secrete IL-17A as schistosomiasis progressed.¹⁷ The results of the present study partially confirmed this: Compared with that in the healthy controls, the level of the IL-17A in serum was significantly lower in the patients with advanced schistosomiasis. Chen et al²² found that $\gamma\delta$ T cells are the main IL-17-producing cells in PBMCs and that IL-17 contributes to granulomatous inflammatory and fibrosing reactions during *S japonica* infection. Wang et al²³




FIGURE 2 Correlation between two subtypes of $\gamma\delta$ T cells and cytokines in patients with advanced schistosomiasis. Correlations among the proportion of the CD27⁺ Vδ1⁺ cells, Vδ1⁺ cells, and the level of cytokines are shown. The cytokines included IL-4, IL-6, IL-10, IL-17A, IL-21, TGF- β , IL-1 β and IL-22. None of the correlations were statistically significant


found that IL-17 concentrations were higher at the acute stage of schistosomiasis compared with that in the other stages. Our findings agreed with the reduction of IL-17 in the advanced stage of infection observed in the mouse model, which might be because the $\gamma\delta$ T cells did not produce any IL-17 in the late stage of infection.¹⁷

There has only been one study of $\gamma\delta$ T cells in patients with schistosomiasis, and that study only included patients with *S. mansoni* and *S. haematobium* infections, not *S. japonicum* infection.¹⁴ They observed changes in the proportion of $\gamma\delta$ T cells among CD3⁺ cells without structural changes of $\gamma\delta$ T cells themselves. A previous study investigated cell functions during infections with other parasites, and showed that circulating $\gamma\delta$ cells were impaired in human chronic infection with cystic echinococcosis.²⁴ $\gamma\delta$ T cells enhanced the expression of *Plasmodium* immunogenic factors and exacerbated subsequent systemic and brain-infiltrating inflammatory $\alpha\beta$ T cell responses.²⁵

Other studies of patients infected with *S. japonicum* tested cytokines in patient serum and showed that IL-21 was increased in acute and chronic infections.^{26,27} However, in our study, there was no significant change in IL-21 levels in the advanced disease group. By contrast, Long et al²⁸ found that *S. japonicum*-induced fibrotic liver tissue had higher IL-13 expression than normal liver tissue. However, IL-13 is not secreted by $\gamma\delta$ T cells, but by traditional $\alpha\beta$ T cells. Li et al²⁹ found that the mRNA levels of *TGFB1* (TGF- β 1) correlated with the stage of fibrosis in patients with chronic disease; however, in the present study, there was no significant difference in serum TGF- β levels between patients with advanced schistosomiasis and the healthy controls. Perhaps, it is necessary to design another experiment, including three groups, including patients with chronic and advanced schistosomiasis, to determine the role of TGF- β . The importance of IL-10 in protecting against pathogen-induced tissue and liver injury has been demonstrated in other infectious disease



FIGURE 3 Correlations among circulating $\gamma\delta$ T cells, cytokines, and the degree of liver fibrosis in patients with advanced schistosomiasis. (A) Correlation between the proportion of the subtypes of $\gamma\delta$ T cells and the degree of liver fibrosis. The subtypes included the CD27⁺ V δ 1⁺ cells, CD27⁻ V δ 1⁻ cells, CD27⁺ V δ 1⁻ cells, CD27⁻ V δ 1⁺ cells, V δ 1⁺ cells, and CD27⁻ cells. (B) Correlation between the level of the cytokines in the serum and the degree of liver fibrosis. The cytokines included IL-4, IL-6, IL-10, IL-17A, IL-21, TGF- β , IL-1 β and IL-22. None of the correlations were statistically significant






FIGURE 4 Correlations between other subtypes of $\gamma\delta$ T cells and cytokines in patients with advanced schistosomiasis. The subtypes included the $CD27^- V\delta1^+$ cells, $CD27^- V\delta1^-$ cells, $CD27^+ V\delta1^-$ cells and $CD27^-$ cells. Levels of IL-17A in serum correlated significantly with the number of $CD27^- V\delta1^-$ $\gamma\delta$ T cells. *: $P < .05$

models³⁰; however, the levels of IL-10 were similar between the two groups in the present study.

In the patients with advanced schistosomiasis, the proportion of the $CD27^- V\delta1^-$ subset among total $\gamma\delta$ T cells correlated positively with IL-17A levels in serum. The $V\delta3$ subtype cells are a very small proportion; therefore, if we consider the $V\delta1^-$ cells as $V\delta2$ cells, then this association suggests that the decline in IL-17A is most likely caused by a decrease in the ratio of $V\delta2$ cells or their activity, which implied that $V\delta2$ cells are the main source of IL-17A in the serum of patients with advanced schistosomiasis. Interestingly, $V\delta2$ cells are the main source of IL-17 in many other infections, such as by *Mycobacterium tuberculosis*³¹ and human immunodeficiency virus.³² $V\delta2$ T cells produce proinflammatory cytokines and chemokines, kill infected cells, secrete growth factors for epithelial cells, and present antigens to $\alpha\beta$ T cells.³³ Besides, although we considered $V\delta1^-$ cells to be $V\delta2$ cells, the specific subtype of $\gamma\delta$ T cells involved and their phenotypic characteristics require further in vitro and in vivo study. Further detailed classification and functional verification of the γ T cells are required. In addition, whether they play a role in the pathological process of schistosomiasis, and the potential mechanism, requires further exploration.

In patients with advanced schistosomiasis, the degree of liver fibrosis increased significantly compared with that in the healthy controls. The increase of HA and PC-III in peripheral blood was related to hepatic fibrosis, which was consistent with our expectations.³⁴ However, in our study, we observed no direct link between the ratios of $\gamma\delta$ T cell subsets and liver fibrosis. It is possible that $\gamma\delta$ T cells only assist with the promotion or mitigation of fibrosis. Another possibility is that other classification methods might need to be explored to reveal the relationship between $\gamma\delta$ T cells and liver fibrosis. Activated $V\gamma9^+$ T cells could act as anti-fibrotic mediators in systemic sclerosis, although decreased responsiveness to isopentenyl

pyrophosphate might play a role in the pathological fibrosis of this disease.¹⁹ In the late stage of schistosomiasis, there was no significant association between hepatic fibrosis and IL-17A producing $\gamma\delta$ T cells, possibly because the body was in an immunosuppressive state, and the effector cells, including $\alpha\beta$ cells, were losing their regulatory ability. However, although IL-17A levels were different between the two groups, there was no correlation between IL-17A and the degree of liver fibrosis. HSCs trigger robust IL-17A production by $\gamma\delta$ T cells through the production of IL-1 β and IL-23 at the early stages of liver fibrosis.³⁵ However, the in vivo function of cytokines that affect HSCs in vitro remains to be determined. Although we did not find any significant relationship between cytokines and fibrosis, this research still serves as a basis for future studies to discover other cytokines associated with liver fibrosis.

Several limitations need to be noted regarding the present study. First, the number of patients was low because there were no new cases of infection by *S japonicum* in Hubei Province for 5 years, and epidemiologically, most cities in this region have been in a state of transmission blocking. Therefore, there were insufficient new cases to fully describe the progress of schistosomiasis and to determine changes in cytokines during the disease process. Secondly, it would be helpful to stain all $\gamma\delta$ T cell subsets specifically, not only staining for one subset and assuming that the remaining cells mostly represent the other major subset ($V\delta2$ cells). The limited experimental conditions meant that we did not obtain intercellular staining of $\gamma\delta$ T cell for IL-17 secretion. Further in vitro experiments should be performed to demonstrate the relationship between $\gamma\delta$ T cells and cytokines.

It is recommended that further research be undertaken in the following areas: First, studies including acute and chronic infection groups are needed. Second, research is also required to identify more cytokines and their intrinsic link to liver fibrosis. Third, the

group of patients need to be enlarged to determine functions of $\gamma\delta$ T cells more clearly, such as the possible correlation of IL-6 and IL-21 with CD27⁺V δ 1⁺ cells.

5 | CONCLUSION

A study of $\gamma\delta$ T cells was carried out in patients with advanced schistosomiasis. The proportion of V δ 1⁺ cells and CD27⁺V δ 1⁺ cells among all $\gamma\delta$ T cells increased. However, there was no significant correlation between the two subsets and the levels of cytokines, including IL-4, IL-6, IL-10, IL-17A, IL-21, TGF- β , IL-1 β or IL-22. In addition, we found no relationship between the level of liver fibrosis and cytokines or subsets of $\gamma\delta$ T cells. Moreover, we found that the serum level of IL-17A correlated positively with the proportion of CD27⁺V δ 1⁺ cells. The present research increased our understanding of the role of $\gamma\delta$ T cells in the immune environment of human schistosomiasis.

ACKNOWLEDGEMENTS

We thank Professor Shao-long Zheng from the Institute for the Prevention and Control of Schistosomiasis in Qian-jiang City for guiding the research, and Qi-lin He from Schistosomiasis Control Hospital in Qian-jiang City for supporting our work.

CONFLICT OF INTEREST

We declare that we have no financial and personal relationships with other people or organizations that can inappropriately influence our work, there is no professional or other personal interest of any nature or kind in any product, service and/or company that could be construed as influencing the position presented in, or the review of, the manuscript.

AUTHOR CONTRIBUTIONS

Li Zheng, Xiaorong Zhou and Jianping Cao: conceived and designed the experiments. Li Zheng: performed the experiments. Jia Yi, Lun Wan, Lixia Wang and Xiaorong Zhou: involved in sample collection. Jianping Cao and Li Zheng: contributed reagents and materials. Li Zheng, Yuan Hu and Yujuan Shen: analysed the data. Li Zheng and Jianping Cao: wrote the paper.

PEER REVIEW

The peer review history for this article is available at <https://publons.com/publon/10.1111/pim.12871>

DATA AVAILABILITY STATEMENT

All datasets generated and analysed during the current study are included in the article. Original raw data are available from the corresponding author upon reasonable request.

ORCID

Li Zheng  <https://orcid.org/0000-0002-1126-0428>

Jianping Cao  <https://orcid.org/0000-0002-1974-0047>

REFERENCES

- Freer JB, Bourke CD, Durhuus GH, Kjetland EF, Prendergast AJ. Schistosomiasis in the first 1000 days. *Lancet Infect Dis*. 2018;18(6):e193-e203. [https://doi.org/10.1016/S1473-3099\(17\)30490-5](https://doi.org/10.1016/S1473-3099(17)30490-5)
- Bluestone JA, Khattry R, Sciammas R, Sperling AI. TCRgammadelta Cells: A specialized T-cell subset in the immune system. *Ann Rev Cell Dev Biol*. 1995;11(1):307-353. <https://doi.org/10.1146/annurev.cb.11.110195.001515>
- Ribot JC, DeBarros A, Pang DJ, et al. CD27 is a thymic determinant of the balance between interferon-gamma- and interleukin 17-producing gammadelta T cell subsets. *Nat Immunol*. 2009;10(4):427-436. <https://doi.org/10.1038/ni.1717>
- Wu YL, Ding YP, Tanaka Y, et al. $\gamma\delta$ T cells and their potential for immunotherapy. *Int J Biol Sci*. 2014;10(2):119-135. <https://doi.org/10.7150/ijbs.7823>
- Kabelitz D, Glatzel A, Wesch D. Antigen recognition by human $\gamma\delta$ T lymphocytes. *Int Arch Allergy Immunol*. 2000;122(1):1-7. <https://doi.org/10.1159/000024353>
- Poles MA, Barsoum S, Yu W, et al. Human immunodeficiency virus type 1 induces persistent changes in mucosal and blood $\gamma\delta$ T Cells despite suppressive therapy. *J Virol*. 2003;77(19):10456-10467. <https://doi.org/10.1128/jvi.77.19.10456-10467.2003>
- Fenoglio D, Poggi A, Catellani S, et al. Vdelta1 T lymphocytes producing IFN-gamma and IL-17 are expanded in HIV-1-infected patients and respond to Candida albicans. *Blood*. 2009;113(26):6611-6618. <https://doi.org/10.1182/blood-2009-01-198028>
- Rakasz E, MacDougall AV, Zayas MT, et al. $\gamma\delta$ T cell receptor repertoire in blood and colonic mucosa of rhesus macaques. *J Med Primatol*. 2000;29(6):387-396. <https://doi.org/10.1111/j.1600-0684.2000.290602.x>
- Biswas P, Ferrarini M. A matter of life and death: More members of the TNF receptor family join human $\gamma\delta$ T lymphocytes. *Eur J Immunol*. 2012;42(3):803-804. <https://doi.org/10.1002/eji.201141983>
- Mangan BA, Dunne MR, O'Reilly VP, et al. Cutting edge: CD1d restriction and Th1/Th2/Th17 cytokine secretion by human V δ 3 T cells. *J Immunol*. 2013;191(1):30-34. <https://doi.org/10.4049/jimmunol.1300121>
- Sandor M, Sperling AI, Cook GA, Weinstock JV, Lynch RG, Bluestone JA. Two waves of gamma delta T cells expressing different V delta genes are recruited into schistosome-induced liver granulomas. *J Immunol*. 1995;155(1):275-284. PMID: 7602105.
- Yu X, Luo X, Xie H, et al. Characteristics of $\gamma\delta$ T cells in *Schistosoma japonicum*-infected mouse mesenteric lymph nodes. *Parasitol Res*. 2014;113(9):3393-3401. <https://doi.org/10.1007/s00436-014-4004-8>
- Xie H, Chen D, Li L, et al. Immune response of $\gamma\delta$ T cells in *Schistosoma japonicum*-infected C57BL/6 mouse liver. *Parasite Immunol*. 2014;36(12):658-667. <https://doi.org/10.1111/pim.12135>
- Schwartz E, Rosenthal E, Bank I. Gamma delta t cells in non-immune patients during primary schistosomal infection. *Immun Inflamm Dis*. 2014;2:56-61. <https://doi.org/10.1002/iid3.18>
- Von Lichtenberg F, Erickson DG, Sadun EH. Comparative histopathology of schistosome granulomas in the hamster. *Am J Pathol*. 1973;72(2):149-178. PMID:4737396.
- Wu W, Feng A, Huang Y. Research and control of advanced schistosomiasis japonica in China. *Parasitol Res*. 2015;114(1):17-27. <https://doi.org/10.1007/s00436-014-4225-x>
- Zheng L, Hu Y, Wang Y, et al. Recruitment of neutrophils mediated by V γ 2 $\gamma\delta$ T cells deteriorates liver fibrosis induced by *Schistosoma japonicum* infection in C57BL/6 mice. *Infect Immun*. 2017;85(8):1-16. <https://doi.org/10.1128/IAI.01020-16>
- Hammerich L, Bangen JM, Govaere O, et al. Chemokine receptor CCR6-dependent accumulation of $\gamma\delta$ T cells in injured liver restricts

- hepatic inflammation and fibrosis. *Hepatology (Baltimore, MD)*. 2014;59(2):630-642. <https://doi.org/10.1002/hep.26697>
19. Markovits N, Bendersky A, Loebstein R, Brusel M, Kessler E, Bank I. Anti-fibrotic characteristics of V γ 9+ $\gamma\delta$ T cells in systemic sclerosis. *Clin Exp Rheumatol*. 2016;34(5):23-29. PMID:26886502.
 20. Kühl AA, Pawlowski NN, Grollich K, et al. Human peripheral $\gamma\delta$ T cells possess regulatory potential. *Immunology*. 2009;128(4):580-588. <https://doi.org/10.1111/j.1365-2567.2009.03162.x>
 21. Davey MS, Willcox CR, Joyce SP, et al. Clonal selection in the human V δ 1 T cell repertoire indicates $\gamma\delta$ TCR-dependent adaptive immune surveillance. *Nat Commun*. 2017;8(1):14760. <https://doi.org/10.1038/ncomms14760>
 22. Chen D, Luo X, Xie H, Gao Z, Fang H, Huang J. Characteristics of IL-17 induction by *Schistosoma japonicum* infection in C57BL/6 mouse liver. *Immunology*. 2013;139(4):523-532. <https://doi.org/10.1111/imm.12105>
 23. Wang X, Fu Q, Song R. Antinuclear antibodies and interleukin responses in patients with *Schistosoma japonicum* infection. *Parasite Immunology*. 2018;40(10):e12577 <http://dx.doi.org/10.1111/pim.12577>
 24. Wang H, Li M, Zhang X, He F, Zhang S, Zhao J. Impairment of peripheral V δ 2 T cells in human cystic echinococcosis. *Experimental Parasitology*. 2017;174:17-24. <http://dx.doi.org/10.1016/j.exppara.2017.01.005>
 25. Ribot JC, Neres R, Zuzarte-Luís V, et al. $\gamma\delta$ -T cells promote IFN- γ -dependent *Plasmodium* pathogenesis upon liver-stage infection. *Proc Natl Acad Sci U S A*. 2019;116(20):9979-9988. <https://doi.org/10.1073/pnas.1814440116>
 26. Zhang Y, Jiang Y, Wang Y, et al. Higher frequency of circulating PD-1high CXCR5⁺CD4⁺ Tfh cells in patients with chronic schistosomiasis. *Int J Biol Sci*. 2015;11(9):1049-1055. <https://doi.org/10.7150/ijbs.12023>
 27. Zhang Y, Wang Y, Jiang Y, et al. T follicular helper cells in patients with acute schistosomiasis. *Parasit Vectors*. 2016;9(1):321. <https://doi.org/10.1186/s13071-016-1602-6>
 28. Long X, Chen Q, Zhao J, et al. An IL-13 promoter polymorphism associated with liver fibrosis in patients with *Schistosoma japonicum*. *PLoS One*. 2015;10:e0135360(8). <https://doi.org/10.1371/journal.pone.0135360>
 29. Li L, Wu T, Huang J, et al. Expression of heat shock protein 47, transforming growth factor-beta 1, and connective tissue growth factor in liver tissue of patients with *Schistosoma japonicum*-induced hepatic fibrosis. *Parasitology*. 2014;48(2):341-351. <https://doi.org/10.1017/S0031182014001115>
 30. Bliss SK, Alcaraz A, Appleton JA. IL-10 prevents liver necrosis during murine infection with *Trichinella spiralis*. *J Immunol*. 2003;171(6):3142-3147. <https://doi.org/10.4049/jimmunol.171.6.3142>
 31. Kulicke CA, Lewinsohn DA, Lewinsohn DM. Clonal enrichments of V δ 2- $\gamma\delta$ T cells in *Mycobacterium tuberculosis*-infected human lungs. *J Clin Invest*. 2020;130(1):68-70. <https://doi.org/10.1172/JCI133119>
 32. Chevalier MF, Bhatnagar N, Didier C, et al. $\gamma\delta$ T-cell subsets in HIV controllers: Potential role of T $\gamma\delta$ 17 cells in the regulation of chronic immune activation. *AIDS*. 2019;33(8):1283-1292. <https://doi.org/10.1097/QAD.0000000000002196>
 33. Morita CT, Jin C, Sarikonda G, Wang H. Nonpeptide antigens, presentation mechanisms, and immunological memory of human V γ 2V δ 2 T cells: discriminating friend from foe through the recognition of prenyl pyrophosphate antigens. *Immunol Rev*. 2007;215(1):59-76. <https://doi.org/10.1111/j.1600-065X.2006.00479.x>
 34. Bin XS, Yao JL, Zheng RQ, Peng XM, Gao ZL. Serum hyaluronic acid, procollagen type III and IV in histological diagnosis of liver fibrosis. *Hepatobiliary Pancreat Dis Int*. 2003;2(1):69-72. PMID:14607650.
 35. Seo W, Eun HS, Kim SY, et al. Exosome-mediated activation of toll-like receptor 3 in stellate cells stimulates interleukin-17 production by $\gamma\delta$ T cells in liver fibrosis. *Hepatology*. 2016;64(2):616-631. <https://doi.org/10.1002/hep.28644>

How to cite this article: Zheng L, Wang L, Hu Y, et al. Higher frequency of circulating V δ 1 $\gamma\delta$ T cells in patients with advanced schistosomiasis. *Parasite Immunol*. 2021;00:e12871. <https://doi.org/10.1111/pim.12871>